

A survey of asymptotic classes and measurable structures

Richard Elwes,
School of Mathematics,
University of Leeds,
Leeds LS2 9JT, England

Dugald Macpherson,
School of Mathematics,
University of Leeds,
Leeds LS2 9JT, England,
h.d.macpherson@leeds.ac.uk

March 24, 2006

1 Introduction

In this article we survey a body of results about classes of finite first order structures in which definable sets have a rather uniform asymptotic behaviour. Non-principal ultraproducts of such classes may have unstable theory, but will be supersimple of finite rank. In addition, they inherit a definable *measure* on definable sets.

The starting point for this work was the following result of [10] on definability in finite fields.

Theorem 1.1 ([10]) *Let $\varphi(\bar{x}, \bar{y})$ be a formula in the language L_{rings} for rings, with $\bar{x} = (x_1, \dots, x_n)$ and $\bar{y} = (y_1, \dots, y_m)$. Then there is a positive constant C , and a finite set D of pairs (d, μ) with $d \in \{0, \dots, n\}$ and μ a non-negative rational number, such that for each finite field \mathbb{F}_q and $\bar{a} \in \mathbb{F}_q^m$,*

$$|\varphi(\mathbb{F}_q^n, \bar{a})| - \mu q^d \leq Cq^{d-(1/2)} \quad (*)$$

for some $(d, \mu) \in D$.

Furthermore, for each $(d, \mu) \in D$, there is a formula $\varphi_{(d, \mu)}(\bar{x})$ which defines in each finite field \mathbb{F}_q the set of tuples \bar{a} such that $(*)$ holds.

This result rests on the Lang-Weil estimates for the number of \mathbb{F} -rational points of an absolutely irreducible variety defined over the finite field \mathbb{F} . The proof uses partial quantifier elimination for pseudofinite fields, derivable from the paper of Ax [2] which introduced pseudofinite fields: any formula $\varphi(\bar{x})$ is a boolean combination of formulas $\exists y(g(\bar{x}, y) = 0)$, where $g \in \mathbb{Z}[\bar{X}, Y]$. (In fact, by adding a set C of constants to the language, the authors arrange that φ is equivalent to a *conjunction* of similar formulas, rather than an arbitrary boolean combination; there is an alternative effective proof in [20], based on Galois stratification.) The reason why μ may be a fraction (rather than an integer as in Lang-Weil), is that the existential quantifier ranges over a finite set. Oversimplifying slightly, the set defined by $\exists y(g(\bar{x}, y) = 0)$ is, for some k , the image of a k -to-1 projection map of the variety defined by $g(\bar{x}, y)$; thus, its measure is $\frac{1}{k}$ times that of $\{(\bar{x}, y) : g(\bar{x}, y) = 0\}$.

As a short application of Theorem 1.1, noted in [10], it follows that the finite field \mathbb{F}_q is not uniformly definable in \mathbb{F}_{q^2} . Theorem 1.1 also yields that pseudofinite fields are supersimple of rank 1, with an associated definable and finitely additive notion of measure on the definable sets.

The present article is a survey of recent work stimulated by Theorem 1.1. The idea, initiated in [40], is to consider classes of finite structures in which definable sets satisfy the same kind of asymptotic behaviour as for finite fields. In the initial work of [40], only a 1-dimensional version was considered, and the error terms were as in Theorem 1.1. More recently, Elwes [18] has developed a theory of higher dimensional asymptotic classes where the error terms are weaker – see Definition 2.1 below.

There is also a notion of *measurable structure*: a supersimple structure of finite SU-rank equipped with the kind of measure function on definable sets which exists, by virtue of Theorem 1.1, in pseudofinite fields. Any ultraproduct of members of an asymptotic class will be measurable (Theorem 3.8 below) but there are also measurable structures which do not even have the finite model property, so cannot arise from asymptotic classes; see Theorem 3.11. Familiar supersimple structures such as the random graph, smoothly approximable structures, and pseudofinite fields, are all measurable.

One of the main results in this area is a theorem of Ryten [48], stemming from earlier work of Ryten and Tomašić [49], that for any natural number d , the collection of all finite simple groups of Lie rank at most d forms an asymptotic class (Theorem 6.1 below). This opens the possibility of variations of the Algebraicity Conjecture of Cherlin and Zilber that any simple group of finite Morley rank is an algebraic group. Namely, the following seems reasonable, and, unlike the Algebraicity Conjecture, incorporates the classes of *twisted* finite simple groups.

Conjecture 1.2 *If G is a simple group with measurable theory, then G is a Chevalley group (possibly of twisted type) over a pseudofinite field.*

In Section 2 below we give the current definition of *asymptotic class*, and some examples. Measure is introduced in Section 3, again with examples and discussion of the connection with asymptotic classes, and with Hrushovski’s notion of *unimodular* theory. Smoothly approximable structures provide an important class of examples, and we describe in Section 4 how these fit into the framework. In Section 5, we survey results from [48] and [49] on measure in ACFA, and an extension of Theorem 1.1 to finite difference fields. This yields the above result on finite simple groups of fixed Lie rank, sketched in Section 6. We turn in Section 7 to measurable groups, and asymptotic classes of groups, of low dimension. The main results here are that any 2-dimensional asymptotic class of groups consists of groups with a uniformly definable soluble subgroup of bounded index (Theorem 7.5), and an analogue of Hrushovski’s result on groups of finite Morley rank in a stable theory which act transitively on a strongly minimal set. The paper concludes with a section on open questions.

If \mathcal{C} is a class of finite structures in a language L , then the *asymptotic theory* of \mathcal{C} is the collection of all sentences which hold in all but finitely many members of \mathcal{C} . Equivalently, it consists of those sentences which hold of any non-principal ultraproduct of members of \mathcal{C} . Notation is introduced locally, but we try to stick to the convention that for a formula $\varphi(\bar{x}, \bar{y})$, $\bar{x} = (x_1, \dots, x_n)$, $\bar{y} = (y_1, \dots, y_m)$, with \bar{y} the parameter variables. The algebraic closure of a field K is denoted \bar{K} . We write $A \downarrow_C B$ to mean that the sets A and B are independent over C (in the sense of model-theoretic non-forking).

Though this material is close to stability and simplicity, our intention is that little knowledge of simplicity theory is needed to follow this paper; the reader can refer to [45] and [57]. For details of the ranks mentioned (S_1 -rank, D -rank, SU-rank), see [34]. For background on the model theory of finite and pseudofinite fields, we refer to the original paper [2], the survey [13], or to [21]. For background on ACFA, see [11]. For the structure of the finite simple groups of Lie type, see [9].

2 Asymptotic classes

Definition 2.1 (Elwes, [18]) Let $N \in \mathbb{N}$, and let \mathcal{C} be a class of finite L -structures, where L is a finite language. Then we say that \mathcal{C} is an N -**dimensional asymptotic class** if the following hold.

(i) For every L -formula $\varphi(\bar{x}, \bar{y})$ where $l(\bar{x}) = n$ and $l(\bar{y}) = m$, there is a finite set of pairs $D \subseteq (\{0, \dots, Nn\} \times \mathbb{R}^{>0}) \cup \{(0, 0)\}$ and for each $(d, \mu) \in D$ a collection $\Phi_{(d, \mu)}$ of pairs of the form (M, \bar{a}) where $M \in \mathcal{C}$ and $\bar{a} \in M^m$, so that $\{\Phi_{(d, \mu)} : (d, \mu) \in D\}$ is a partition of $\{(M, \bar{a}) : M \in \mathcal{C}, \bar{a} \in M^m\}$, and

$$|\varphi(M^n, \bar{a})| - \mu|M|^{\frac{d}{N}} = o(|M|^{\frac{d}{N}})$$

as $|M| \rightarrow \infty$ and $(M, \bar{a}) \in \Phi_{(d, \mu)}$.

(ii) Each $\Phi_{(d, \mu)}$ is \emptyset -definable, that is to say $\{\bar{a} \in M^m : (M, \bar{a}) \in \Phi_{(d, \mu)}\}$ is uniformly \emptyset -definable across \mathcal{C} .

We may write D_φ for D , and will call $\{\Phi_{(d, \mu)} : (d, \mu) \in D\}$ a (*definable*) *asymptotic partition*. We write $h(\varphi(M^n, \bar{a})) := (\text{Dim}(\varphi(M^n, \bar{a})), \text{Meas}(\varphi(M^n, \bar{a}))) := (d, \mu)$ where $(M, \bar{a}) \in \Phi_{(d, \mu)}$, except that if $d = \mu = 0$ we work with the convention that $\text{Dim}(\varphi(M^n, \bar{a})) = -1$. We call \mathcal{C} a *weak asymptotic class* when \mathcal{C} satisfies the asymptotic criteria (i) for all φ , but the $\Phi_{(d, \mu)}$ are not assumed to be definable.

Remark 2.2 1. In this context the o -notation in (i) means the following: for every $\varepsilon > 0$ there is $Q \in \mathbb{N}$ such that for all $M \in \mathcal{C}$ with $|M| > Q$ and all $\bar{a} \in M^m$ where $(M, \bar{a}) \in \Phi_{d, \mu}$, we have

$$|\varphi(M^n, \bar{a})| - \mu|M|^{\frac{d}{N}} < \varepsilon|M|^{\frac{d}{N}}.$$

2. Unlike the presentation in [40] and [18], the definition covers all formulas $\varphi(\bar{x}, \bar{y})$, not just those in which \bar{x} is a singleton. It is a theorem (Theorem 2.3 below) that this is equivalent to the same condition just for formulas $\varphi(x, \bar{y})$.

3. The focus of [40] was on the case when $N = 1$ (the case for finite fields).

4. The general theory below works equally well with tighter error terms: replace $o(|M|^{d/N})$ by $C|M|^{\frac{d}{N} - \frac{1}{2N}}$, where C is a constant. An advantage from adopting the weaker error term above is that for smoothly approximable structures M , even if M is not unidimensional, M is the direct limit of a sequence of ‘envelopes’ which forms an asymptotic class. With the tighter error terms, this appears unclear. For example, if p, q are distinct primes, and M is the disjoint union of two infinite-dimensional vector spaces, one over \mathbb{F}_p , the other over \mathbb{F}_q , is M approximated by an asymptotic class satisfying the tighter error terms?

5. It is easy to create artificial examples (just using sets with a unary predicate) in which the μ are irrational, or even transcendental. We do not know of natural examples with transcendental μ .

6. It is immediate that any reduct of an asymptotic class is a weak asymptotic class. However, the definability clause (ii) can be lost under reducts. Elwes [18, Section 2.2] has shown that if \mathcal{C} is an asymptotic class, and \mathcal{D} is a class of finite structures uniformly interpretable in \mathcal{C} , then \mathcal{D} is a weak asymptotic class. If \mathcal{C} and \mathcal{D} are uniformly bi-interpretable (without parameters), then \mathcal{D} will also be an asymptotic class. This is relevant to Section 6 below (bi-interpretations between classes of simple groups and fields or difference fields).

The following theorem gives a more easily recognised criterion for being an asymptotic class. By ensuring that the condition is really one on one-variable definable sets, it gives an analogy to o -minimality, and related minimality notions.

Theorem 2.3 (Lemma 2.1.2 of [18]) *Suppose that \mathcal{C} is a class of finite structures which satisfies Definition 2.1 (clauses (i) and (ii)) for $n = 1$, i.e. for definable sets in 1 variable. Then \mathcal{C} is an N -dimensional asymptotic class.*

Sketch proof. The proof is inductive. For a definable subset X of M^{n+1} consider the projection $\pi : M^{n+1} \rightarrow M$ to the first variable, and apply the assumption to $\pi(X)$, and the inductive hypothesis to the fibres X_a ($a \in \pi(X)$). The definability clause (ii) is essential.

We draw attention to the following *non-example*. The class of all finite total orders is not an asymptotic class of any dimension; for, as a varies through a finite ordering, the formula $x < a$ defines an arbitrary proportion of the domain. In fact, by Proposition 3.8 below, any nonprincipal ultraproduct of an asymptotic class is supersimple, so cannot interpret a partial order with an infinite chain (i.e. cannot have the *strict order property*).

Example 2.4 1. By Theorem 1.1, the collection of all finite fields forms a 1-dimensional asymptotic class. This class, when restricted to fixed characteristic, has an important expansion. For any fixed p , and positive integers m, n with $m \geq 1$, $n > 1$, and $(m, n) = 1$, there is a 1-dimensional asymptotic class $\mathcal{C}_{(m,n,p)}$ of difference fields, namely $\mathcal{C}_{(m,n,p)} = \{(\mathbb{F}_{p^{kn+m}}, \text{Frob}^k) : k > 0\}$ – see Theorem 5.8. The automorphism here is not uniformly definable in the field. The classes $\mathcal{C}_{(m,n,p)}$ are significant for finite simple groups: $\mathcal{C}_{1,2,2}$ is uniformly parameter bi-interpretable (in a sense to be defined) with the classes of Suzuki groups ${}^2B_2(2^{2k+1})$ and the Ree groups ${}^2F_4(2^{2k+1})$, and $\mathcal{C}_{(1,2,3)}$ with the class of Ree groups ${}^2G_2(3^{2k+1})$.

2. It is shown in [40, Theorem 3.14] that the collection of all finite cyclic groups is a 1-dimensional asymptotic class. This follows from the partial quantifier elimination (‘near model completeness’) of Szemielew [52] for abelian groups. The multiplicative groups of finite fields form a subclass.

3. For any odd prime p , the class \mathcal{C}_p of finite extraspecial p -groups of exponent p forms a 1-dimensional asymptotic class (see [40, Proposition 3.11]). Here, a group G is *extraspecial* if $G' = Z(G) = \Phi(G)$ (the Frattini subgroup), and G' is isomorphic to the cyclic group \mathbb{Z}_p . An extraspecial group of exponent p is just a central product of copies of a specific group of order p^3 . The groups in \mathcal{C}_p are bounded-by-abelian (in fact, \mathbb{Z}_p -by-abelian), but not abelian-by-bounded, that is, they do not have an abelian normal subgroup of bounded index. The quotient $G/Z(G)$ carries definably the structure of a vector space over the field \mathbb{F}_p , equipped with an alternating bilinear form which comes from the commutator map. An infinite ultraproduct of groups in \mathcal{C}_p will be ω -categorical, smoothly approximable (see Section 4), and supersimple of rank 1, but not stable. See also Theorem 7.3, and the remark after Lemma 7.4, for partial converses.

4. If q is a prime power with $q \equiv 1 \pmod{4}$, then there is a graph P_q , a *Paley graph*, with vertex set the finite field \mathbb{F}_q , with vertices a, b joined if $a - b$ is a square. By [40], the class \mathcal{C} of all Paley graphs is a 1-dimensional asymptotic class. Essentially, the reason is that by a theorem of Bollobás and Thomason [6] (see also [7, Ch.XIII.2]), if U, W are disjoint sets of vertices of P_q with $m := |U \cup W|$, and $v(U, W)$ is the number of vertices of P_q joined to everything in U and to nothing in W , then

$$|v(U, W) - 2^{-m}q| \leq \frac{1}{2}(m - 2 + 2^{-m+1})q^{\frac{1}{2}} + m/2.$$

It follows that an infinite ultraproduct of Paley graphs is elementarily equivalent to the random graph, so has quantifier elimination. This persists in sufficiently large finite Paley graphs, so corresponding asymptotic estimates hold for *all* formulas in one variable, and hence, by Theorem 2.3, for all formulas. The definability clause of Definition 2.1 is easily verified.

There are analogous classes, interpretable in finite fields, associated with other homogeneous structures. For example, if one considers primes $q \equiv 3 \pmod{4}$, and defines $a \rightarrow b$ (on \mathbb{F}_q) whenever $a - b$ is a square, one obtains the *Paley tournaments*.

These approximate the random tournament as above; the analogue of the Bollobás-Thomason result was proved in [22]. For more on this, see [53], and the results of Tomašić [55] mentioned at the end of Section 3.

There is an analogue of the Paley graphs for arity 3 (so a class of 3-hypergraphs), considered in [40, Example 3.6]. For $q \equiv 1 \pmod{4}$ one considers a hypergraph on \mathbb{F}_q where $\{a, b, c\}$ is an edge if a, b, c are distinct and $(a-b)(b-c)(a-c)$ is a square. The asymptotic theory of such hypergraphs is that of the generic homogeneous *two-graph* (a 3-hypergraph which is a reduct of the random graph), not the generic 3-hypergraph.

5. By Proposition 3.3.2 of [18], if \mathcal{M} is a smoothly approximable structure, then \mathcal{M} is the union of a chain of ‘envelopes’ which form an asymptotic class. See Section 4.

6. Suppose that \mathcal{C} is a class of finite structures such that every infinite ultraproduct of members of \mathcal{C} is strongly minimal. Then \mathcal{C} is a 1-dimensional asymptotic class (Lemma 2.5 of [40]). This is an easy consequence of Theorem 2.3.

In particular, let d be a positive integer, and let \mathcal{C}_d be the collection of all finite graphs of valency d whose automorphism group is transitive on the vertex set. Then \mathcal{C}_d is a 1-dimensional asymptotic class. The reason is that any infinite ultraproduct M of members of \mathcal{C}_d is itself vertex transitive: take the ultraproduct of 2-sorted structures, with a second sort for the automorphism group. It follows that M is strongly minimal, so the last paragraph applies.

In the next section, it will be shown that if \mathcal{C} is an N -dimensional asymptotic class, then any infinite ultraproduct of members of \mathcal{C} is supersimple of rank at most N (and, furthermore, *measurable*). A number of other model-theoretic properties of ultraproducts can be recognised directly from asymptotics. One such is the following.

Proposition 2.5 ([40]) *Let \mathcal{C} be a 1-dimensional asymptotic class. Then some ultraproduct of members is unstable if and only if there is a formula $\varphi(x, \bar{y})$, and for each $k \in \mathbb{N}$ some $M \in \mathcal{C}$ and $\bar{a}_1, \dots, \bar{a}_k \in M^{\ell(\bar{y})}$ with*

- (a) $|\varphi(M, \bar{a}_i)| \geq k$ for each $i = 1, \dots, k$, and
- (b) $|\varphi(M, \bar{a}_i) \Delta \varphi(M, \bar{a}_j)| \geq k$ for all distinct $i, j \in \{1, \dots, k\}$.

In Chapter 5 of [18], finitary criteria are given for ultraproducts to be ω -categorical, or all to be elementarily equivalent, or all to be 1-based.

3 Measurable structures

It was already noted in [10, 4.10, 4.11] that, because of Theorem 1.1, pseudofinite fields are supersimple of S_1 -rank 1, and that there is a definable ‘measure’ on the definable sets. Below, following Section 5 of [40], we generalise this.

Definition 3.1 An infinite L -structure M is *measurable* if there is a function $h : \text{Def}(M) \rightarrow \mathbb{N} \times \mathbb{R} \cup \{(0, 0)\}$ (we write $h(X) = (\text{Dim}(X), \text{Meas}(X)) = (\text{Dim}, \text{Meas})(X)$) such that the following hold.

1. For each L -formula $\varphi(\bar{x}, \bar{y})$ there is a finite set $D \subset \mathbb{N} \times \mathbb{R}^{>0} \cup \{(0, 0)\}$, so that for all $\bar{a} \in M^m$ we have $h(\varphi(M^n, \bar{a})) \in D$.
2. If $\varphi(M^n, \bar{a})$ is finite then $h(\varphi(M^n, \bar{a})) = (0, |\varphi(M^n, \bar{a})|)$.
3. For every L -formula $\varphi(\bar{x}, \bar{y})$ and all $(d, \mu) \in D_\varphi$, the set $\{\bar{a} \in M^m : h(\varphi(M^n, \bar{a})) = (d, \mu)\}$ is \emptyset -definable.

4. (Fubini) Let $X, Y \in \text{Def}(M)$ and $f : X \rightarrow Y$ be a definable surjection. Then there are $r \in \omega$ and $(d_1, \mu_1), \dots, (d_r, \mu_r) \in (\mathbb{N} \times \mathbb{R}^{>0}) \cup \{(0, 0)\}$ so that if $Y_i := \{\bar{y} \in Y : h(f^{-1}(\bar{y})) = (d_i, \mu_i)\}$, then $Y = Y_1 \cup \dots \cup Y_r$ is a partition of Y into non-empty disjoint definable sets. Let $h(Y_i) = (e_i, \nu_i)$ for $i \in \{1, \dots, r\}$. Also let $c := \text{Max}\{d_1 + e_1, \dots, d_r + e_r\}$, and suppose (without loss) that this maximum is attained by $d_1 + e_1, \dots, d_s + e_s$. Then $h(X) = (c, \mu_1 \nu_1 + \dots + \mu_s \nu_s)$.

If $X \in \text{Def}(M)$ and $h(X) = (d, \mu)$, we call d the *dimension* of X and μ the *measure* of X , and h the *measuring function*. We say that a complete theory T is *measurable* if it has a measurable model (see Remark 3.7 (1)).

Example 3.2 The basic motivating example of a measurable structure is a pseudofinite field. Essentially, this was shown in [10]. If $X \subset F^n$ is definable (F a pseudofinite field) then $\text{Dim}(X)$ (which equals its S_1 -rank or D -rank) is just the algebraic-geometric dimension of the Zariski closure of X in \tilde{F}^n . The measure of any absolutely irreducible variety in F will be 1. Measurability of any pseudofinite field was used in [27], in a new proof of the well-known fact that any almost simple algebraic group has a universal cover. The key point was that, by measure considerations, if F is a pseudofinite field and G_1, G are connected algebraic groups defined over F , and $f : G_1 \rightarrow G$ is an isogeny defined over F , then $|\text{Ker}(f) \cap G_1(F)| = |G(F) : f(G_1(F))|$.

A key question is whether, conversely, every measurable field is pseudofinite (see Section 8). An algebraically closed field cannot be measurable – see the remark after Proposition 3.14.

Generalising the case of pseudofinite fields, Proposition 3.8 below shows that asymptotic classes yield measurable structures.

Definition 3.1 is slightly different from that in [40], since we do not specify in the definition that M has supersimple theory. However, it follows from the next few lemmas, in particular Corollary 3.6, that indeed if M is measurable, then $\text{Th}(M)$ is supersimple of finite rank. Similar arguments were communicated to the authors by Ryten, and can also be found in [31]. The computations below are with D -rank, but by [34, Section 6], in a supersimple theory in which D -rank is finite, D -rank, S_1 -rank, and SU -rank all agree for formulas. See [34] for definitions of these ranks.

Lemma 3.3 *Let $n \in \omega$ and let A_1, \dots, A_n be definable sets, where $\text{Dim}(A_i) = d$ for each i .*

(i) $\text{Dim}(\bigcup_{i=1}^n A_i) = d$.

(ii) *If, in addition we have $\text{Dim}(A_{i_1} \cap A_{i_2}) < d$ for each distinct i_1, i_2 , then*

$$(\text{Dim}, \text{Meas})\left(\bigcup_{i=1}^n A_i\right) = \left(d, \sum_{i=1}^n \text{Meas}(A_i)\right).$$

Proof. We proceed by induction on n . For $n = 2$, pick any distinct $a_1, a_2, a_3 \in M$, and define $f : A_1 \cup A_2 \rightarrow \{a_1, a_2, a_3\}$ by $f(\bar{x}) := a_1$ if $\bar{x} \in A_1 \setminus A_2$; $f(\bar{x}) := a_2$ if $\bar{x} \in A_2 \setminus A_1$; and $f(\bar{x}) = a_3$ if $\bar{x} \in A_1 \cap A_2$. Then (i) and (ii) are immediate from the Fubini condition.

Suppose now that both statements hold for $n = k - 1$. Then given A_1, \dots, A_k , we may first apply the inductive hypothesis to A_1, \dots, A_{k-1} , and derive statement (i) by applying the case $n = 2$ to $\bigcup_{i=1}^{k-1} A_i$ and A_k .

Now for (ii), we know by the inductive hypothesis that $(\text{Dim}, \text{Meas})\left(\bigcup_{i=1}^{k-1} A_i\right) = \left(d, \sum_{i=1}^{k-1} \text{Meas}(A_i)\right)$. Now $A_k \cap \bigcup_{i=1}^{k-1} A_i = \bigcup_{i=1}^{k-1} (A_k \cap A_i)$, and so by (i) we know that $\text{Dim}(A_k \cap \bigcup_{i=1}^{k-1} A_i) < d$. Therefore we may apply the case $n = 2$ to find that

$\text{Meas}(\bigcup_{i=1}^k A_i) = \text{Meas}(\bigcup_{i=1}^{k-1} A_i) + \text{Meas}(A_k) = \sum_{i=1}^{k-1} \text{Meas}(A_i) + \text{Meas}(A_k) = \sum_{i=1}^k \text{Meas}(A_i)$, as required. \square

Corollary 3.4 *(Dim, Meas) is monotonic, that is whenever $A \subseteq B$ are definable, then $(\text{Dim}, \text{Meas})(A) \leq (\text{Dim}, \text{Meas})(B)$ (under the lexicographic ordering).*

For convenience we assume that all the following occurs in the home sort.

Lemma 3.5 *Let X be a definable set, $\varphi(\bar{x}, \bar{y})$ an L -formula, and $(\bar{b}_i : i \in \omega)$ an indiscernible sequence where for each $i \in \omega$ we have $\varphi(M^n, \bar{b}_i) \subseteq X$. Suppose that $\{\varphi(M^n, \bar{b}_i) : i \in \omega\}$ is inconsistent. Then $\text{Dim}(X) > \text{Dim}(\varphi(M^n, \bar{b}_i))$.*

Proof. Suppose not. Suppose $\text{Dim}(X) = \text{Dim}(\varphi(M^n, \bar{b}_i)) = d$. Then, as $\{\varphi(M^n, \bar{b}_i) : i \in \omega\}$ is inconsistent, by compactness there exists some minimal k such that $\text{Dim}(\varphi(M^n, \bar{b}_1) \cap \dots \cap \varphi(M^n, \bar{b}_{k+1}) \cap \varphi(M^n, \bar{b}_{k+2})) < d$.

For $i \geq 1$ define $A_i := \varphi(M^n, \bar{b}_1) \cap \dots \cap \varphi(M^n, \bar{b}_k) \cap \varphi(M^n, \bar{b}_{k+i})$.

Notice that by indiscernibility and the minimality of k , we have $\text{Dim}(A_i) = d$, and for $i_1 \neq i_2$ also $\text{Dim}(A_{i_1} \cap A_{i_2}) < d$. Say $\text{Meas}(A_i) = \mu$. Thus by Lemma 3.3, for any $t \geq 1$ we have $(\text{Dim}, \text{Meas})(\bigcup_{i=1}^t A_i) = (d, t\mu)$. But then by Corollary 3.4 $\text{Meas}(X) \geq t\mu$ for all t , which is clearly impossible. \square

Corollary 3.6 *For any definable set X , we have $D(X) \leq \text{Dim}(X)$.*

Proof. We proceed by induction on r , showing (*) that if $\text{Dim}(X) \leq r$ then $D(X) \geq r$. The case $r = 0$ is automatic from clause (2) of Definition 3.1.

For the inductive step, assume (*) below r , and suppose for a contradiction that $\text{Dim}(X) = r$, and $D(X) \geq r + 1$.

By definition of D -rank, there is an indiscernible sequence $(\bar{b}_i : i \in \omega)$ and an L -formula $\varphi(\bar{x}, \bar{y})$ where $\{\varphi(\bar{x}, \bar{b}_i) : i \in \omega\}$ is inconsistent, and for each $i \in \omega$ we have $D(\varphi(M^n, \bar{b}_i)) \geq r$ and $\varphi(M^n, \bar{b}_i) \subseteq X$. By Lemma 3.5, we have $\text{Dim}(\varphi(M^n, \bar{b}_i)) < r$ for each i . It follows by the inductive hypothesis that $D(\varphi(M^n, \bar{b}_i)) < r$ for each i , which is a contradiction. \square

Remark 3.7 1. It is immediate from the definition, and noted in [40], that measurability is a property of a *theory*; that is, if M is measurable and $M \equiv N$, then N is measurable.

2. Less obviously, if M is measurable, and we adjoin to M finitely many sorts from M^{eq} , then the resulting structure is measurable – see [40, Proposition 5.10]. In fact, the class of measurable theories is closed under bi-interpretability.

3. A measurable structure may have many different measuring functions. For example, for the random graph, there is a measure corresponding to any edge probability p , where $0 < p < 1$: we let p be the measure of the set of neighbours of a vertex [40, 5.12]. This is generalised in Theorem 3.9 below, and the remarks after it. If a theory is measurable, then, rather as with Theorem 2.3, the measuring function is determined by its restriction to definable sets in one variable.

4. It can happen that in a measurable structure M there are definable sets X_1 and X_2 with $\text{Dim}(X_1) = \text{Dim}(X_2)$ but $D(X_1) \neq D(X_2)$. For example, consider a structure (M, P, E) where P is a unary predicate picking out an infinite subset with infinite complement, and E is an equivalence relation partitioning $P(M)$ into infinitely many infinite classes, but with $\neg P(M)$ a single E -class. The $D(P(M)) = 2$ and $D(\neg P(M)) = 1$, but we can artificially choose a dimension and measure such that $\text{Dim}(P(M)) = \text{Dim}(\neg P(M)) = 2$.

One can modify this example to produce a measurable structure in which D -rank is not definable. Indeed, consider the structure $(M, E, F_i)_{i \in \omega}$, where E is an equivalence relation whose classes $\{X_i : i \in \mathbb{Z}\}$ are all infinite, and for each $i \in \omega$,

F_i is an equivalence relation which agrees with E except on X_i , which it partitions into infinitely many infinite classes. There is a measuring function on M under which each X_i has dimension 2. But $D(X_i) = 1$ for $i < 0$ and $D(X_i) = 2$ for $i \geq 0$, and the latter are not uniformly definable.

5. If M is measurable with measuring function $(\text{Dim}, \text{Meas})$, and $S \subset M^n$ is definable, there is an induced finitely additive probability measure μ_S on the σ -algebra generated by the definable subsets of S : for definable $X \subset S$, put

$$\mu_S(X) = \begin{cases} \frac{\text{Meas}(X)}{\text{Meas}(S)} & \text{if } \text{Dim}(X) = \text{Dim}(S) \\ 0 & \text{if } \text{Dim}(X) < \text{Dim}(S) \end{cases}$$

If S is defined by the formula $\psi(\bar{x})$, we sometimes write it as μ_ψ .

6. Ben-Yaacov [3] defines, in a measurable theory, the relation $\bar{a} \downarrow_C^d \bar{b}$, to mean $\text{Dim}(\bar{a}/C) = \text{Dim}(\bar{a}/C \cup \{\bar{b}\})$, where, if $\bar{a} = (a_1, \dots, a_n)$,

$$\text{Dim}(\bar{a}/C) := \min\{\text{Dim}(\varphi(M^n/\bar{c})) : \varphi(\bar{a}, \bar{c}), \bar{c} \text{ from } C\}.$$

He shows that this relation coincides with non-dividing.

Proposition 3.8 ([18]) *Let $\mathcal{C} = \{M_i : i \in \omega\}$ be an N -dimensional asymptotic class, and $M = \prod M_i$ be an infinite ultraproduct of members of \mathcal{C} . Then M is measurable, with $(\text{Dim}, \text{Meas})(M) = (N, 1)$.*

Proof. Use the definability clause (ii) of Definition 2.1 to define $(\text{Dim}, \text{Meas})(\varphi(\bar{x}, \bar{a}))$ for any $\bar{a} \in M^m$. Easy asymptotic arguments show that this is indeed a measure. \square

It follows that all the classes of examples listed in Examples 2.4 yield corresponding examples of measurable structures. In particular, the random graph is measurable, as it is an ultraproduct of Paley graphs; and any infinite vertex transitive graph of finite valency is measurable.

We mention some further examples and constructions. The first comes from the generic predicate construction of [12].

Theorem 3.9 (5.11 of [40]) *Let T be a complete measurable theory over a language L with quantifier elimination, eliminating the quantifier \exists^∞ , such that for all $M \models T$ and $A \subset M$, $\text{acl}(A) = \text{dcl}(A)$. Let P be a unary predicate not in L , let $L' := L \cup \{P\}$, let S be a sort of T , and let $T_{P,S}$ be the model companion of the theory of L' -structures satisfying T , with P interpreted by a subset of S . Let T' be any completion of $T_{P,S}$. Then T' is measurable.*

In the proof, one uses the measuring function μ_T for T , fixes $p \in \mathbb{R}$ with $0 < p < 1$, and for $M \models T_{P,S}$, assigns to $P(M)$ the measure $p\mu_T(S)$. The partial quantifier elimination of [12, 2.6] makes this measure extendable to all definable sets. We expect that the assumption $\text{acl}(A) = \text{dcl}(A)$ is not needed, but it was assumed in [40] to avoid complications.

As noted in [40], it follows from Theorem 3.9 that for any $k \geq 2$, the universal homogeneous k -uniform hypergraph Γ_k (that is, a universal homogeneous structure with a single symmetric irreflexive k -ary relation) is measurable. It is not obvious that for $k \geq 2$ this structure is an ultraproduct of an asymptotic class, as there is no obvious analogue for the Paley graphs of Example 2.4 (4). However, Beyarslan [personal communication] has shown that for any $k \geq 2$, Γ_k is interpretable in a pseudofinite field, so it must at least be an ultraproduct of a weak asymptotic class.

The motivating example of a measurable structure is a pseudofinite field. Any pseudofinite field arises as the fixed field of the automorphism in a generic difference

field, that is, in a model of ACFA. This, and the last theorem, suggests the following construction technique for measurable structures, given by Hrushovski in Proposition 11.1 of [29]. The definability of measure was not explicitly stated in [29], but uniqueness was, and as noted in [18, 3.4.2], definability follows from uniqueness by Beth's Theorem.

Theorem 3.10 ([29]) *Let D be any strongly minimal set over a language L , assume that $\text{Th}(D)$ has the definable multiplicity property (DMP) and elimination of imaginaries, let σ be a generic automorphism of a sufficiently saturated model of $\text{Th}(D)$, and let $K = \text{Fix}(\sigma)$, an L -substructure. Then K is measurable, of dimension 1.*

We remark that under assumption (DMP), by [12, 3.11(2)], there is a generic automorphism (i.e. a model companion of the theory of expansions of D by automorphisms).

Recall Hrushovski's fusion construction [30], which, given two strongly minimal sets M_1, M_2 with (DMP) in disjoint languages L_1, L_2 respectively, yields a new strongly minimal set M in $L_1 \cup L_2$ whose reduct to each L_i is M_i . It is shown in [30] that this construction preserves the (DMP). Thus, there is a generic automorphism σ of M , and $\text{Fix}(\sigma)$ will be measurable. As shown in Section 3.4 of [18], this yields the following, when applied to the fusion of two algebraically closed fields of characteristics p_1 and p_2 .

Theorem 3.11 ([18]) *Let L_1, L_2 be disjoint languages for rings, and $L := L_1 \cup L_2$, and p_1, p_2 be distinct primes. Then there is a measurable structure M whose reduct to each language L_i is a pseudofinite field of characteristic p_i .*

Note that M is not elementarily equivalent to an ultraproduct of an asymptotic class, since there do not exist positive integers a_1, a_2 with $p_1^{a_1} = p_2^{a_2}$.

Example 3.12 ([40]) It is shown in [40] that for any field F , any infinite vector space over F , in the language of F -modules, is a measurable structure. In the particular case when F is infinite, such a structure cannot be an ultraproduct of an asymptotic class. The point here, essentially, is that if M is *any* strongly minimal set with (DMP) and definable Skolem functions, then M is measurable, with dimension and measure equal to Morley rank and degree. In vector spaces over an infinite field, this is applied after naming a non-zero vector by a constant symbol.

The following definition is given in [25].

Definition 3.13 Let T be a complete theory. We say that T is *unimodular* if for any $M \models T$, definable sets X, Y in M^{eq} , and definable surjections $f_i : X \rightarrow Y$ such that f_i is k_i -to-1 (for $i = 1, 2$, and with k_i a positive integer), we have $k_1 = k_2$.

We say that M is unimodular if $\text{Th}(M)$ is.

Clearly any theory with the finite model property (e.g. an ultraproduct of an asymptotic class) is unimodular. In fact, the following is almost immediate from the Fubini condition.

Proposition 3.14 *Let M be measurable. Then M is unimodular.*

In applications of measure seen by the authors so far, only unimodularity (plus finite rank supersimplicity with definability of some dimension or rank) is used.

It follows immediately from Proposition 3.14 that an algebraically closed field cannot be measurable. For example, in characteristic not equal to 2, if F is an algebraically closed field, then the identity map is 1-to-1, but the map $x \mapsto x^2$ is 2-to-1 on $F \setminus \{0\}$. More generally, we have the following theorem. Recall that a sufficiently saturated structure M with simple theory is said to be *1-based* if, for any subsets A, B of M^{eq} , $A \downarrow_{\text{acl}^{\text{eq}}(A) \cap \text{acl}^{\text{eq}}(B)} B$.

Theorem 3.15 (Theorem 4.2.6 of [18]) *Any unimodular stable theory of finite U -rank (and hence any measurable stable theory) is 1-based.*

The main point in the proof is that by [25] (see also 2.4.15 and 5.3.2 of [45]), every minimal type of the theory will be 1-based. By the coordinatisation of such structures by minimal types, this is sufficient (see 2.5.8 of [45]).

We have the following easy observation.

Proposition 3.16 *Let M be ω -categorical. Then M is unimodular.*

Proof. Suppose that M is ω -categorical, and X, Y are definable sets in M^{eq} and $f_i : X \rightarrow Y$ are definable surjections with f_i k_i -to-1 (for $i = 1, 2$, and with $k_i \in \mathbb{N}$). By adding finitely many sorts to M , naming finitely many parameters, and adding dummy variables, we may suppose that X, Y are disjoint subsets of M , and X, Y and the f_i are \emptyset -definable. Let $a \in X$, and $D = \text{acl}(a) \cap (X \cup Y)$. Then D is finite, by ω -categoricity, and is closed under the f_i and f_i^{-1} . Thus, f_i induces a k_i -to-1 map from $D \cap X$ onto $D \cap Y$ and it follows by counting that $k_1 = k_2$. \square

Itay Ben-Yaacov [3] has investigated the connections between measurable theories and continuous model theory, and in particular, with measure algebras in an unbounded continuous logic. His main result is a measure-theoretic extension of the independence theorem [34] for simple theories. A version was proved earlier by very different means, for pseudofinite fields, by Tomašić [56].

We mention also some work of Tomašić [55] which, in the context of pseudofinite fields, makes connections between measure and exponential sums, and the independence theorem. It generalises the use of characters in the proof of Example 2.4 (4) for Paley graphs. See also the survey article by Szönyi [53] for combinatorial applications of related results.

First, consider a measurable structure M with measuring function (d, μ) , and a definable set S in M , with induced probability measure μ_S on the definable subsets of S , yielding a measure space $(S, \mathcal{M}_S, \mu_S)$. Then if $f : S \rightarrow \mathbb{C}$ is measurable, we may form the integral $\int_S f d\mu_S$. Tomašić investigates additive and multiplicative characters on a pseudofinite field F , and proves the following. As elsewhere in this paper, \tilde{F} denotes the algebraic closure of the field F .

Theorem 3.17 ([55]) *Let X be an absolutely irreducible variety over a pseudofinite field F and let f be a regular function on X (or a rational function when X is irreducible). Suppose either*

(i) χ is a multiplicative character of F of order $k > 1$ and f is not a k^{th} power of a rational function on \tilde{X} (the corresponding variety over \tilde{F}), or

(ii) χ is a nontrivial additive character of F of the form $\chi_{a,1}$ (notation from [55]) and f is not of the form $g^p - g$ for any $g \in \tilde{F}(\tilde{X})$.

Then $\int_X \chi \circ f = 0$.

The author then considers reducts of a pseudofinite field F of the following form. Let X be an absolutely irreducible variety over F , let f be a regular function on $X \times X$, and let $\chi : X \rightarrow \mathbb{C}$ be a multiplicative character of order $k > 1$ (with the assumption of Theorem 3.17(i)). Given $x, y \in X(F)$, define the binary relation $R_j(x, y)$ to hold if $\chi(f(x, y)) = e^{2\pi i j/k}$. Then there is a binary structure $(X(F), R_0, \dots, R_{k-1})$ interpretable in F . Ultraproducts of Paley graphs are really a special case, with $X(F) = F$, $f(x, y) = x - y$, and χ the quadratic character. Using Theorem 3.17, Tomašić shows that under certain conditions such reducts are ω -categorical, and their measures, and the corresponding structures in finite fields, share the ‘equidistribution’ properties of the theorem of Bollobás and Thomason in Example 2.4 (4). In particular, if $X(F) = F$, f is a symmetric polynomial defining a conic, and χ is a quadratic character, then (F, R_0) is ω -categorical.

4 Smoothly approximable structures

The class of *smoothly approximable* structures is a class of ω -categorical supersimple structures of finite rank which properly contains the class of ω -categorical ω -stable structures (so in particular the totally categorical structures). A deep structure theory is developed in [14], which includes, for example, proofs of the equivalence of smooth approximation, Lie coordinatisability, and other notions, such as ‘strong 4-quasifiniteness’. In [14] there is also a proof of a version of quasi-finite axiomatisability, a Lachlan-style shrinking and stretching theory, results on definable groups (they must be finite-by-abelian-by-finite) and much else.

Following Definition 2.1.1 of [14], we say that a finite substructure N of a structure M is a *k-homogeneous substructure* of M if all \emptyset -definable relations on M induce \emptyset -definable relations of N , and for any pair \bar{a}, \bar{b} of k -tuples from N , they have the same type in N if and only if they have the same type in M . An ω -categorical structure M is *smoothly approximated* if it is a union of a chain $(M_i : i \in \omega)$ of finite substructures, where for each i , M_i is an $|M_i|$ -homogeneous substructure of M .

Smoothly approximated structures with primitive automorphism groups were classified in [32]. Based on this, a list of rank 1 *Lie geometries* is identified in [14, Section 2.1.2]. Typical examples are vector spaces (or their projective and affine versions) over finite fields, possibly equipped with bilinear forms, but there is a rather more mysterious example, the quadratic geometry. The sorts and languages to handle these geometries are chosen with care in [14] to ensure flexibility in handling slightly different automorphism groups (e.g. semilinear automorphisms, which involve field automorphisms), and quantifier elimination and weak elimination of imaginaries in certain cases. The authors define a *Lie coordinatised structure* to be one built by covering constructions from Lie geometries in a way indexed by a tree of finite height, and then prove that a countable structure is smoothly approximated if and only if it is Lie coordinatisable (i.e. bi-interpretable with a Lie coordinatised structure).

It is easily seen that the Lie geometries arise as direct limits of 1-dimensional asymptotic classes. For example, given a finite field \mathbb{F}_q , let \mathcal{C} be the collection of all finite dimensional vector spaces V over \mathbb{F}_q , equipped with a non-degenerate alternating bilinear form $\beta : V \times V \rightarrow \mathbb{F}_q$. We may code β by introducing a binary relation symbol R_a for each $a \in \mathbb{F}_q$, with $R_a(x, y)$ whenever $\beta(x, y) = a$. Witt’s Lemma, which says that partial isometries extend to total isometries, gives quantifier elimination, so we can reduce to considering the cardinalities of sets of the form $\{x : \beta(x, v_1) = a_1 \wedge \dots \wedge \beta(x, v_r) = a_r\}$, where $a_1, \dots, a_r \in \mathbb{F}_q$, and $v_1, \dots, v_r \in V$ are linearly independent. Such a set has size $\frac{1}{q^r}|V|$. Thus, there is some resemblance with the way, in the random graph, 1-types over a finite set $\{x_1, \dots, x_r\}$ are determined by edges/non-edges to each x_i , with all sets of adjacencies equiprobable (assuming edge probability $\frac{1}{2}$); but unlike in the Paley graphs, we get precise results on sizes of definable sets. The random graph is not smoothly approximable.

Theorem 4.1 (Section 3.3.2 of [18]) (i) *Let M be a smoothly approximable structure. Then M is the union of a sequence of substructures $(M_i : i < \omega)$ such that each M_i is a $|M_i|$ -homogeneous substructure of M , and $\{M_i : i \in \omega\}$ is an asymptotic class.*

(ii) *Every smoothly approximable structure is measurable.*

The M_i are also called *envelopes* in [14]. Theorem 4.1(i) rests on rather precise information in [14, Proposition 5.2.2] on the sizes of definable sets in envelopes; these cardinalities are given by polynomials in certain dimensions. As a result, for smoothly approximated structures for which only one canonical projective geometry

is involved in the coordinatisation, the asymptotics for the envelopes are much better than that required in Definition 2.1 – see [18, Proposition 3.3.5].

Part (ii) of Theorem 4.1 follows immediately from (i) and Proposition 3.8, since any smoothly approximable structure satisfies the asymptotic theory of any approximating chain $(M_i : i \in \omega)$ of envelopes.

One question throughout this paper concerns the extent to which the examples of asymptotic classes and measurable structures go beyond finite and pseudofinite fields, and structures interpretable in them. It is conceivable that any smoothly approximable structure is interpretable in a product of pseudofinite fields, in which case Theorem 4.1 does not provide new examples. Certainly, we have

Proposition 4.2 ([40]) *Let M be a smoothly approximable Lie geometry. Then M is interpretable in a pseudofinite field.*

As an example, consider an \aleph_0 -dimensional vector space V over a finite field \mathbb{F}_p , equipped with a non-degenerate symmetric bilinear form $\beta : V \times V \rightarrow \mathbb{F}_p$. This is approximated by the family $(V_n : n \geq 1)$, where V_n is an n -dimensional vector space. We may identify V_n with \mathbb{F}_p^n , viewed as a vector space over \mathbb{F}_p . There is a trace map $\text{Tr} : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$, and we may put $\beta(x, y) = \text{Tr}(xy)$ for any $x, y \in \mathbb{F}_p^n$. The trace map is uniformly definable (so definable in the limit). Indeed, its kernel has index p in $(\mathbb{F}_p^n, +)$, and is uniformly defined as $\{x^p - x : x \in \mathbb{F}_p^n\}$ (Hilbert’s Theorem 90). To define the trace, we just specify its value on each of the p cosets of the kernel.

5 Measure and difference fields

The families of Suzuki and Ree finite simple groups are not uniformly interpretable in finite fields. However, as is clear from their constructions, they are uniformly interpretable in certain finite difference fields. In this and the next section, we describe work of Ryten, based on joint work of Ryten and Tomašić, which ensures that *all* families of finite simple groups of fixed Lie rank form asymptotic classes.

Ryten’s starting point is the following theorem of Hrushovski.

Theorem 5.1 ([30]) *Let \tilde{K} be an algebraically closed field of characteristic p , let $n \in \mathbb{N}$, and $q = p^t$. Suppose $V(\tilde{x})$ is an algebraic variety over \tilde{K} , and $W(\tilde{x}\tilde{z}) \subset V(\tilde{x}) \times V^q(\tilde{z})$ is an irreducible subvariety, where the polynomials defining V^q are the images under the automorphism $y \mapsto y^q$ of those defining V . Assume $\text{Dim}(W) = \text{Dim}(V) = d$, and that the projection $\pi_1 : W \rightarrow V$ is dominant of degree δ and $\pi_2 : W \rightarrow V^q$ is quasi-finite of purely inseparable degree δ' . Then there is a constant C depending on the total degree of W such that*

$$|\{|\tilde{x}\tilde{z} \in W(\tilde{K}) : \tilde{z} = \tilde{x}^q\}| - \frac{\delta}{\delta'} q^d| \leq C q^{d-\frac{1}{2}}.$$

From this, Hrushovski proves

Theorem 5.2 ([30]) *Any non-principal ultraproduct of pairs $(\tilde{\mathbb{F}}_p, \text{Frob}^k)$ is a model of ACFA.*

Using these results, Ryten and Tomašić prove the following.

Theorem 5.3 ([49]) *Let $\theta(\tilde{x}, \tilde{y})$ be a formula in the language of difference rings. Then there is a constant $C \in \mathbb{R}^+$ and a finite set D of pairs (d, μ) with $d \in \mathbb{Z} \cup \{\infty\}$ and $\mu \in \mathbb{Q}^+ \cup \{\infty\}$ such that in each difference field $(\tilde{\mathbb{F}}_p, \text{Frob}^k)$, and for any $\tilde{a} \in \tilde{\mathbb{F}}_p^m$,*

$$|\theta(\tilde{x}, \tilde{a}) - \mu p^{kd}| \leq C p^{k(d-\frac{1}{2})}$$

holds for some $(d, \mu) \in D$.

Moreover, for each $(d, \mu) \in D$ there is a formula $\theta_{d, \mu}(\bar{y})$ such that for each $(\tilde{\mathbb{F}}_p, \text{Frob}^k)$, the above estimate holds for $\theta(\tilde{\mathbb{F}}_p, \bar{a})$ with (d, μ) if and only if $(\tilde{\mathbb{F}}_p, \text{Frob}^k) \models \theta_{d, \mu}(\bar{a})$.

The proof has a somewhat different presentation in Chapter 2 of [48]. The theorem is derived from Theorem 5.1 rather as Theorem 1.1 is derived from the Lang-Weil estimates. Formulas with quantifiers are handled using Theorem 5.2, together with the following partial quantifier elimination for ACFA.

Proposition 5.4 (1.5 and 1.6 of [11]) *If $\theta(\bar{x}, \bar{y})$ is a formula in the language of difference rings, then*

$$ACFA \models \theta(\bar{x}, \bar{y}) \Leftrightarrow \bigvee_{i=1}^k \exists t \theta_i(\bar{x}, \bar{y}, t),$$

where $\theta_i = \theta_i(\bar{x}, \sigma(\bar{x}), \dots, \sigma^\ell(\bar{x}), \bar{y}, \sigma(\bar{y}), \dots, \sigma^\ell(\bar{y}), t, \sigma(t), \dots, \sigma^\ell(t))$ is a quantifier free formula in the language of rings, and for any $(M, \sigma) \models ACFA$ and $(\bar{x}_0, \bar{y}_0, t_0) \in \theta_i(M)$, t_0 is algebraic (in the sense of fields) over $\{\sigma^i(\bar{x}_0), \sigma^i(\bar{y}_0) : 0 \leq i \leq \ell\}$.

Ryten and Tomašić also note that, by Theorem 5.3, in a model of ACFA, the family of *finite dimensional* sets is measurable, under a natural variant of Definition 3.1.

In [48], Ryten uses Theorem 5.3 to investigate an important class of finite difference fields. Fix a prime p , and integers m, n with $m \geq 1$, $n > 1$, and $(m, n) = 1$. Let $\mathcal{C}_{m, n, p}$ be the collection $\{(\mathbb{F}_{p^{kn+m}}, \text{Frob}^k) : k > 0\}$ of finite difference fields. Observe that Frob^k is a solution in σ of the equation $\text{Frob}^m \circ \sigma^n = \text{id}$.

Consider the following conditions on a difference field (K, σ) . Below, with m, n as above, if (K, σ) is a difference field of characteristic p with $\text{Frob}^m \circ \sigma^n = \text{id}$, we say that the extension of difference fields $(K, \sigma) \subseteq (L, \sigma')$ is *generic* if $\text{Fix}(\text{Frob}^m \circ \sigma'^n) = K$.

- (1) K is a pseudofinite field of characteristic p .
- (2) σ is an automorphism of K which satisfies $\text{Frob}^m \circ \sigma^n = \text{id}$.
- (3) Suppose $U = U(x_{11} \dots x_{n1} \dots x_{1N} \dots x_{nN})$ is an absolutely irreducible variety defined over K and $\sigma(U) = \sigma(U)(y_{11} \dots y_{1n} \dots y_{N1} \dots y_{Nn})$, the variety obtained from U by applying σ to the coefficients of the defining polynomials. Suppose $V \subset U \times \sigma(U)$ is an absolutely irreducible variety defined over K , whose definition includes the equations $y_{ij} = x_{i+1, j}$ and $y_{nj}^{p^m} = x_{1j}$ for $i = 1, \dots, n-1$ and $j = 1, \dots, N$. Suppose that V projects generically onto U and $\sigma(U)$, and that W is a K -algebraic set properly contained in V . Then there is a point $x \in V(K) \setminus W(K)$ such that $x = (a, b)$, where $a = (a_{ij} : 1 \leq i \leq n, 1 \leq j \leq N)$, $b = (b_{ij} : 1 \leq i \leq n, 1 \leq j \leq N)$, $a \in U$, $b \in \sigma(U)$, and $b_{ij} = \sigma(a_{ij})$ for each i, j .

- (4) Let $K \subseteq L \subseteq H$ be a tower of finite field extensions. Suppose that $(K, \sigma) \subseteq (L, \sigma')$ is an extension of difference fields which is generic over K . Then there is an extension of difference fields $(L, \sigma') \subseteq (H, \sigma'')$ such that (H, σ'') is a generic extension of (K, σ) .

It can be shown that (1)–(4) are first order expressible.

Proposition 5.5 ([48]) *Let (M, τ) be a model of ACFA of characteristic p , let $K := \text{Fix}(\text{Frob}^m \circ \tau^n)$, and put $\sigma := \tau|_K$. Then $(K, \sigma) \models (1) - (4)$.*

For conditions (1)–(3), only basic facts about ACFA are needed. The proof that $(K, \sigma) \models (4)$ uses Theorem 5.2, together with the existence of generic extensions of automorphisms for finite extensions of fields, proved by Ryten.

By Proposition 5.5, conditions (1)–(4) axiomatise a theory, denoted $\text{PSF}_{(m,n,p)}$. Ryten proves for $\text{PSF}_{(m,n,p)}$ results similar to those known for ACFA and PSF. The completions (and types) are described through the following theorem.

Theorem 5.6 ([48]) *Let (F, σ) and (E, τ) be models of $\text{PSF}_{(m,n,p)}$ with a common substructure K (so $\sigma|_K = \tau|_K$). Then $(F, \sigma) \equiv_K (E, \tau)$ if and only if $(F \cap \tilde{K}, \sigma|_{F \cap \tilde{K}}) \cong_K (E \cap \tilde{K}, \tau|_{E \cap \tilde{K}})$.*

From this, near model completeness is proved: that is, for every formula $\varphi(\bar{x})$ there is a formula $\theta(\bar{x})$ which is a boolean combination of existential formulas (in fact, formulas of a rather specific type), such that $\text{PSF}_{(m,n,p)} \vdash \varphi(\bar{x}) \leftrightarrow \theta(\bar{x})$.

Using Theorem 5.2, Ryten proves the following theorem.

Theorem 5.7 ([48]) *$\text{PSF}_{(m,n,p)}$ is the asymptotic theory of the class $\mathcal{C}_{(m,n,p)}$ of finite difference fields.*

We show part of this, that every non-principal ultraproduct

$$(N, \sigma) = \Pi_{i \in \mathbb{N}} (\mathbb{F}_p^{n k_i + m}, \text{Frob}^{k_i}) / \mathcal{U}$$

satisfies $\text{PSF}_{(m,n,p)}$. Indeed, if $(M, \tau) = \Pi_{i \in \mathbb{N}} (\tilde{\mathbb{F}}_p^{n k_i + m}, \text{Frob}^{k_i}) / \mathcal{U}$, then $(M, \tau) \models \text{ACFA}$ by Theorem 5.2. Also, $N = \text{Fix}(\text{Frob}^m \circ \sigma^n)$ and $\sigma = \tau|_N$, so $(N, \sigma) \models \text{PSF}_{(m,n,p)}$ by Proposition 5.5.

Theorem 5.8 ([48]) *$\mathcal{C}_{(m,n,p)}$ is a 1-dimensional asymptotic class.*

To see this, let $\mathcal{D}_p := \{(\tilde{\mathbb{F}}_p, \text{Frob}^k) : k \in \mathbb{N}\}$, a class of difference fields. There is a bijection $\text{Fix} : \mathcal{D}_p \rightarrow \mathcal{C}_{(m,n,p)}$, with $\text{Fix}((\tilde{\mathbb{F}}_p, \text{Frob}^k)) = (\mathbb{F}_p^{k n + m}, \text{Frob}^k)$. There is a corresponding function Fix^{Fix} defined on the set of formulas of the language L_{diff} of difference rings, defined inductively. If $(K, \sigma) \in \mathcal{D}_p$ and $\text{Fix}((K, \sigma) = (M, \sigma) \in \mathcal{C}_{(m,n,p)}$, then for any L_{diff} formula $\varphi(y)$ and tuple \bar{a} from M ,

$$(M, \sigma) \models \varphi(\bar{a}) \Leftrightarrow (K, \sigma) \models \varphi^{\text{Fix}}(\bar{a}).$$

From this and Theorem 5.2, the uniform asymptotic estimates for formulas in $\mathcal{C}_{(m,n,p)}$ are rapidly derived. The definability clause (Definition 2.1(ii)) requires a little more work.

Finally, Ryten shows that in a *pure* pseudofinite field (in fact, in any pure bounded PAC field) the only definable field automorphisms are powers of the Frobenius. Thus, $\text{PSF}_{(m,n,p)}$ is a proper expansion of the theory of pseudofinite fields of characteristic p , and $\mathcal{C}_{(m,n,p)}$ is not interpretable in any class of pure finite fields.

In [48, Chapter 3], Ryten proves a number of other results about $\text{PSF}_{(m,n,p)}$, indicating that its complexity is somewhere between that of PSF and ACFA. As for pseudofinite fields in [10], it is possible to add constants to the language to ensure, in the partial quantifier elimination, that only *positive* boolean combinations of existential formulas are used, so model-completeness in the expanded language is obtained. Model-theoretic independence is characterised algebraically. Elimination of imaginaries (over a language with constants for an elementary submodel) is proved.

6 Asymptotic classes of simple groups

There is a natural notion of *uniform parameter bi-interpretability* between two classes \mathcal{C} and \mathcal{D} of finite structures. We do not give it formally, but it requires a matching between \mathcal{C} and \mathcal{D} , and that each element of \mathcal{C} is bi-interpretability with

the corresponding element of \mathcal{D} , possibly using parameters, but with uniformity of the interpretation across the families.

The non-abelian finite simple groups, excluding the sporadics, are the alternating groups $\text{Alt}(n)$, the classical groups of Lie rank n over the finite field \mathbb{F}_q , and certain exceptional groups, namely $E_6(q), E_7(q), E_8(q), F_4(q), G_2(q)$, and the twisted groups ${}^3D_4(q), {}^2E_6(q)$, the Ree groups ${}^2F_4(2^{2k+1})$, the Suzuki groups ${}^2B_2(2^{2k+1})$, and the Ree groups ${}^2G_2(2^{3k+1})$. We may regard $\text{Alt}(n)$ as having Lie rank n . We emphasise that the twisted groups (including the classical unitary groups) do not arise over algebraically closed fields, since their definition depends on finite field extensions. Hence they do not have finite Morley rank analogues.

Ryten shows that, for every family of non-abelian finite simple groups of fixed Lie rank, other than those of Suzuki and Ree groups, the family is uniformly parameter bi-interpretable with a class of finite fields. It follows that such a family is an asymptotic class. Care is needed here with the definability clause in Definition 2.1, because of the role of parameters in the bi-interpretation.

For the Suzuki and Ree groups, the situation is more complicated. In fact, the classes of groups $\{{}^2F_4(2^{2k+1}) : k \in \mathbb{N}\}$ and $\{{}^2B_2(2^{2k+1}) : k \in \mathbb{N}\}$ are uniformly parameter bi-interpretable with $\mathcal{C}_{(1,2,2)}$, and $\{{}^2G_2(3^{2k+1}) : k \in \mathbb{N}\}$ is uniformly parameter bi-interpretable with $\mathcal{C}_{(1,2,3)}$. A word of explanation is necessary here. The twisted group ${}^2E_6(q)$, for example, is the fixed point set inside $E_6(q^2)$ of a certain automorphism σ which is a product of a graph automorphism (an automorphism arising from the symmetry of the E_6 Dynkin diagram), and the Frobenius. All this data is definable in the field \mathbb{F}_{q^2} . However, in the case ${}^2G_2(3^{2k+1})$, for example, the automorphism σ is a product of a graph automorphism and the automorphism $x \mapsto x^{3^k}$, a proper (unbounded) power of the Frobenius. So to define ${}^2G_2(3^{2k+1})$ uniformly, it is necessary to be able to define the automorphism Frob^k of $\mathbb{F}_{3^{2k+1}}$, and hence to work in $\mathcal{C}_{(m,n,p)}$.

The interpretability of the finite simple groups of fixed Lie rank in the corresponding field or difference field is well-known. In particular, it was known that any non-principal ultraproduct of such a family was interpretable in ACFA, and supersimple of finite rank. The fact that the fields are interpretable in the groups was proved by Simon Thomas [54].

The above results, together with Theorem 5.8, yield the following theorem of Ryten.

Theorem 6.1 ([48]) *Any family of non-abelian finite simple groups of fixed Lie rank is an asymptotic class.*

We mention two further results, beyond the measurable/asymptotic class context. A group is *pseudofinite* if it is an infinite model of the theory of finite groups.

Theorem 6.2 ([41]) *Let G be a stable pseudofinite group. Then G has a definable soluble subgroup of finite index.*

The key point is that a non-principal ultraproduct of finite simple groups of fixed Lie rank, though supersimple, is not stable, as a pseudofinite field is interpretable in it. The proof of the theorem below rests on this, together with slightly delicate arguments with chain conditions, and basic facts about finite nilpotent groups. It is not possible here to strengthen ‘soluble’ to nilpotent: as noted independently by Khelif and Zilber, there is a metabelian pseudofinite stable group which is not nilpotent-by-finite.

Theorem 6.3 (Wilson [58]) *Every pseudofinite simple group is elementarily equivalent to a Chevalley group (possibly twisted) over a pseudofinite field.*

In [47], Point proves that an ultraproduct of simple Chevalley groups of fixed type (possibly twisted) is *isomorphic* to a Chevalley group over the ultraproduct of the fields, and is simple.

7 Groups of low dimension

Theorem 6.1 above yields immediately the converse to Conjecture 1.2: any Chevalley group (possibly twisted) over a pseudofinite field is measurable. In this section we sketch beginnings of a general structure theory for measurable groups, with a view to Conjecture 1.2.

The central definability result for groups of finite Morley rank is the Zilber Indecomposability Theorem, which generalises the Indecomposability Theorem for algebraic groups. In the supersimple case, we have the following, proved first by Hrushovski [29] for S_1 -theories, that is, supersimple theories of finite S_1 -rank in which the S_1 -rank is definable.

Theorem 7.1 (Wagner[57]) *Let G be a definable group in a supersimple theory of finite rank, and let X_i ($i \in I$) be definable subsets of G . Then there is a definable subgroup H of G such that*

- (i) $H \leq \langle X_i : i \in I \rangle$, with each element of H a product of a bounded number of elements of the X_i and X_i^{-1} , and
- (ii) X_i/H is finite for each $i \in I$.

Proof. This follows from 5.4.5 and 5.5.4 of [57], by compactness. \square

This is used repeatedly in arguments described below. It also yields, for example, that any non-abelian definably simple measurable group is simple, and that in a measurable theory, the derived subgroup of a definable group is always definable.

A second useful tool, already heavily used by Wagner for groups in supersimple theories, is Schlichting's Theorem [51], proved independently by Bergman and Lenstra [5]. Two subgroups H and K of a group G are *commensurable* if each of $|H : H \cap K|$ and $|K : H \cap K|$ are finite.

Theorem 7.2 ([51], [5]) *Let G be a group, H a subgroup, and suppose there is $n \in \mathbb{N}$ such that for all $g \in G$, $|H : H \cap H^g| \leq n$. Then there is a normal subgroup of G which is commensurable with H .*

It is well-known that every rank 1 superstable group is abelian-by-finite. In the supersimple case, one expects 'abelian-by-finite' to be replaced by 'finite-by-abelian-by-finite', in view of the extraspecial groups considered in Example 2.4(3). However, this is an open question. The following result of Elwes and Ryten is proved in [18, 6.0.11]; it was proved under the stronger assumption of measurability in [40].

Theorem 7.3 ([18]) *If G is unimodular supersimple of rank 1, then G has a definable normal subgroup H of finite index, such that H has a finite central subgroup Z , with H/Z abelian.*

The proof is a counting argument with unimodularity. It uses the fact that if G is a BFC group (a group with a finite bound on the size of its conjugacy classes) then the derived subgroup G' is finite; see Theorem 3.1 of [44]. The key to the counting argument is the following lemma, proved in [40] under measurability; the proof under unimodularity is easily extracted from that of [18, 6.0.11]. It eliminates the possibility of measurable groups with finitely many non-identity conjugacy classes, all of full dimension.

Lemma 7.4 ([18]) *Let G be a group defined in a supersimple unimodular theory of finite rank. Then there is $g \in G \setminus \{1\}$ such that $C_G(g)$ is infinite.*

In [40], a cruder argument is given to prove the asymptotic class analogue of Theorem 7.3. It states that any 1-dimensional asymptotic class of *finite* groups consists of groups which are bounded-by-abelian-by-bounded. The argument applies to any class of finite groups all of whose ultraproducts are supersimple of rank 1.

These results, and the analogue for superstable groups, suggest that any measurable 2-dimensional group should be soluble-by-finite. This question is open, but some progress in this direction was made in [19]. Using Theorems 7.3 and 7.1 (to replace any 1-dimensional normal subgroup by a definable one), Elwes and Ryten show that any counterexample interprets a simple group G of dimension and S_1 -rank 2. Further information is obtained: for example, the group G must have conjugacy classes (in fact, infinitely many) of dimension 1, and it must also have at least 1, but finitely many, conjugacy classes of dimension 2. Such examples could not arise as ultraproducts of an asymptotic class, by the classification of finite simple groups. Thus, they prove

Theorem 7.5 ([18]) *Let \mathcal{C} be a 2-dimensional asymptotic class of groups. Then there is $d \in \mathbb{N}$ such that each group G in \mathcal{C} has a subgroup of index at most d which is soluble of derived length at most 4 (and uniformly definable in the class).*

Elwes and Ryten have also investigated measurable structures (G, X) , where G is a group with a definable faithful action on X . The intended analogy is with [39], on the structure of primitive permutation groups of finite Morley rank (see also [8] in this volume). More specifically, they generalise the theorem of Hrushovski [24], that in a stable theory, a connected group of finite Morley rank acting definably and transitively on a strongly minimal set must have Morley rank at most 3, and either acts regularly (so is a strongly minimal group) or is of form $\text{AGL}(1, F)$ or $\text{PSL}(2, F)$ in the natural action (F an algebraically closed field).

A permutation group G on a set X is *primitive* if there is no proper non-trivial G -invariant equivalence relation on X , or equivalently if each stabiliser G_x ($x \in X$) is a maximal subgroup of G . If G , X , and the G -action on X are definable in some structure, we say that (G, X) is *definably primitive* if there is no *definable* proper non-trivial G -invariant equivalence relation on X , or equivalently if point stabilisers are definably maximal. Definable primitivity implies transitivity, as the orbit equivalence relation is definable. We shall say that (G, X) is a *measurable group action* if G , X , and an action of G on X are all definable in a measurable structure. It is an *asymptotic group action* if it is elementarily equivalent to an infinite ultraproduct of an asymptotic class of group actions.

Proposition 7.6 ([19]) *Let (G, X) be a measurable faithful group action, with G infinite, and suppose that G acts faithfully and definably primitively on X . If $\text{Dim}(X) < \text{Dim}(G)$, then G is primitive on X .*

The proof uses Theorems 7.2 and 7.1. To start the proof (and illustrate a useful argument), define \sim on X , putting

$$x \sim y \Leftrightarrow |G_x : G_x \cap G_y| < \infty,$$

(where G_x denotes the stabiliser of $x \in X$). Then \sim is a G -invariant equivalence relation on X , so is definable, so either there is a single \sim -class, or \sim -classes are singletons. The first case is easily eliminated. For if there is a single \sim -class, then (essentially because measurable theories eliminate the quantifier \exists^∞), there is a fixed upper bound on the indices $|G_x : G_x \cap G_y|$ over all $x, y \in X$. Thus, by

Theorem 7.2, there is $N \triangleleft G$ uniformly commensurable to all the G_x , and the proof of 7.2 yields that N is definable. Since N is normal and non-trivial, it is transitive on X . However, as N is commensurable with G_x , the orbit of x under N is finite, a contradiction as X is infinite.

Theorem 7.7 ([19]) *Let (G, X) be an asymptotic group action which is faithful and definably primitive, with $\text{Dim}(X) = 1$. Then $\text{Dim}(G) \leq 3$, and one of the following holds.*

(i) $\text{Dim}(G) = 1$, and G has a definable subgroup B of finite index which is torsion-free divisible abelian and acts regularly on X . (In this situation, for the description we only require measurability, not an asymptotic group action.)

(ii) $\text{Dim}(G) = 2$. In this case, there is a definable pseudofinite field K , and a definable infinite subgroup T of the multiplicative subgroup K^* , such that G is isomorphic to $K^+ \cdot T$ (a subgroup of $\text{AGL}(1, K)$) in its natural action on $(K, +)$.

(iii) $\text{Dim}(G) = 3$. Here G has a unique minimal normal subgroup T , which is definable and isomorphic to $\text{PSL}_2(K)$ (K a pseudofinite field) and its action on X is its natural action on the projective line.

A non-regular example in (i) would be $K^+ \rtimes \{+1, -1\}$, where K is a pseudofinite field in characteristic 0. For (ii), one might take K to be an ultraproduct of fields \mathbb{F}_p ($p \equiv 1 \pmod{4}$) and T to be the squares of K .

Sketch Proof. First, if $\text{Dim}(G) = 1$, then G is finite-by-abelian-by-finite by Theorem 7.3, and in particular G has a subgroup B of finite index defined as the union of the finite conjugacy classes of G . As $B \triangleleft G$, B is transitive (the orbits of B would yield a definable G -invariant partition of X). Also the derived subgroup B' is finite, so trivial by definable primitivity, so B is abelian, and hence acts regularly on X . The remaining analysis in this case is easy.

If $\text{Dim}(G) = 2$, then by Theorem 7.5, G is soluble-by-finite. A minimal normal subgroup A of G will be abelian, and transitive on X (by primitivity); as A is abelian it acts regularly on X . It follows that we may identify A with X and G with a semidirect product $A \rtimes G_1$, where G_1 is the stabiliser of the identity of A (in its action by conjugation. For the full description in (ii), a version of the Zilber Field Theorem is used.

Finally, suppose that $\text{Dim}(G) \geq 3$. In this case one may apply the O’Nan-Scott Theorem, the standard reduction theorem for finite primitive permutation groups; see [37] for a careful account of it. This, and the fact that $\text{Dim}(X) = 1$, reduces us to a situation where G has a unique minimal definable normal subgroup T , and T is simple and $T \leq G \leq \text{Aut}(T)$. A further analysis, using extensive information about finite simple groups, reduces to the case when $T = \text{PSL}(2, K)$ (K a pseudofinite field) in its action on the projective line. Tools used include Aschbacher’s description of maximal subgroups of finite classical groups [1].

As with Theorem 7.5, it would be very nice to have a version of Theorem 7.7 for groups with measurable theories, but the above proof makes heavy use of finite group theory.

We mention one further result which can be proved under measure, but is open just under supersimplicity.

Theorem 7.8 ([40]) *Let G be an ω -saturated measurable group. Then G has an infinite abelian subgroup.*

In the proof, one looks at a counterexample G of minimal dimension. The exponent must be finite (as otherwise there are infinite cyclic subgroups), and by the minimality assumption, every definable subgroup is finite or of finite index. Let $N := \{g \in G : g^G \text{ is finite}\}$. Then N is finite or of finite index. But $N \neq \{1\}$, for

by Lemma 7.4 there is $g \in G \setminus \{1\}$ with $C_G(g)$ infinite; then $|G : C_G(g)|$ is finite, so $g \in N$. If N is infinite, then we may assume that $N = G$, so $|G : C_G(g)|$ is finite for all $g \in G$, and it is then easy to construct an infinite abelian subgroup. Finally, if N is finite but non-trivial, apply similar arguments to G/N .

8 Further questions

If M is a 1-dimensional measurable structure, then it has SU-rank 1, and so algebraic closure gives a pregeometry. It is natural to ask whether this pregeometry satisfies the Zilber Trichotomy: trivial, locally modular (so 1-based) non-trivial, and field-like. A trivial geometry arises for the random graph or any vertex transitive graph of finite valency. Smoothly approximable Lie geometries are locally modular, and, except for a pure set, non-trivial. And pseudofinite fields (as well as ultraproducts of classes $\mathcal{C}_{m,n,p}$ and some examples arising from Theorem 3.11) are non locally modular, and interpret a pseudofinite field.

It should be possible to construct a non locally modular 1-dimensional measurable structure which does not interpret an infinite group. Let M be the saturated, strongly minimal set constructed in [26], which is known to have (DMP), is not locally modular, and has no infinite interpretable group. Then M admits a generic automorphism σ . It seems that a variant of Theorem 3.11 for finite Morley rank, mentioned in Remark (viii) at the end of [29], should apply in M^{eq} (to ensure elimination of imaginaries), and yield that if $N = \text{Fix}(\sigma)$ then the induced structure on N is measurable of dimension 1. We can ensure that N is not 1-based: for algebraic closure in N is the same as that induced from M , and (M, id) embeds in some model (M', σ') of the theory of generic automorphisms of models of $\text{Th}(M)$, so that $N := \text{Fix}(\sigma')$ contains M , so contains a witness to non 1-basedness. Also, an infinite definable group in N would yield, by a group configuration argument, an infinite definable group in M , which is impossible.

However, we ask

Question 1. If M is a 1-dimensional measurable structure which is a non-principal ultraproduct of an asymptotic class, and M is not 1-based, must M interpret an infinite field?

The question can also be asked for structures interpretable in PSF. In the same vein, we ask whether measurability can be used to sharpen the group configuration results of [4].

Question 2. Is every ω -categorical measurable structure 1-based?

For this, a key example is Hrushovski's construction in [28] of a ω -categorical non-locally modular supersimple rank 1 structure. It is unimodular, by Proposition 3.16, but we do not know whether it is measurable. There is more chance of a positive answer to Question 2 for structures interpretable in PSF. On Question 2, Elwes [18, 5.2.3] has obtained some partial results analogous to Proposition 8 of [25].

Question 3. Is every measurable field pseudofinite?

By the results of Ax [2], a field F is pseudofinite if and only if it is perfect, quasifinite (i.e. has absolute Galois group $\hat{\mathbb{Z}}$), and satisfies the PAC condition; the latter asserts that any absolutely irreducible variety defined over F has an F -rational point. It is straightforward that any supersimple field of finite rank is perfect. By an argument of Scanlon ([40, Theorem 5.18 and Appendix], see also [50]), any measurable field is quasifinite. We do not know whether every measurable field satisfies the PAC condition. It was shown in [46] that any supersimple division ring is commutative and has absolute Brauer group, so the norm from any finite extension to the field is surjective. By [42], any generic elliptic or hyperelliptic curve

defined over a supersimple field F has an F -rational point. If F is supersimple and has a unique quadratic extension (e.g. if F is measurable), then by [43] *any* elliptic curve defined over F has an F -rational point.

Question 4. Is every measurable simple group a (possibly twisted) Chevalley group over a pseudofinite field?

Of course, Question 4 requires a positive answer to Question 3. Question 4 looks difficult, but seems more feasible (without using the classification of finite simple groups) for simple groups interpretable in models of Ryten’s theories $\text{PSF}_{(m,n,p)}$ of measurable pseudofinite difference fields. We remark that Hrushovski [29, Theorem 9.2] classified simple groups interpretable in bounded PAC fields. They are essentially Chevalley groups (possibly twisted), but include unexpected examples which do not arise over pseudofinite fields. Dello Stritto [personal communication] has partial results which, modulo a positive answer to Question 3, are likely to answer Question 4 positively for groups with a BN pair of rank at least 2.

It would also be interesting to obtain, without use of the classification, structural results on asymptotic classes of finite simple groups.

Question 5. Is every 2-dimensional measurable group soluble-by-finite? Equivalently, is every non-abelian infinite measurable simple group of dimension at least 3?

Following Proposition 4.2, we ask:

Question 6. Is every smoothly approximable structure interpretable in a pseudofinite field?

Finally, it would be helpful to compare measurability with some similar notions in the literature. One such is unimodularity, and we ask

Question 7. Is there a supersimple finite rank unimodular theory which is not measurable?

An example to try would again be Hrushovski’s construction from [28]. In Question 7 one might replace ‘unimodular’ by the stronger condition ‘has the finite model property’.

Question 8. (Macintyre) Is there a natural example of a measurable structure in which some measures are transcendental numbers?

We finish with some comments on other counting and measure principles.

There is a notion of measure investigated by Keisler in [33], and more recently in [31]. The latter consider a sort X in a sufficiently saturated model M , and define a Keisler measure on X to be a finitely additive probability measure on $\text{Def}_1(X)$, the collection of parameter-definable subsets of X . A type, for example, is just a zero-one measure. Here we are not requiring a measure on subsets of all powers of M , so there is no Fubini-like assumption. If M is measurable, then for any sort S , there is a definable Keisler measure on S : the two measures will agree for definable subsets of S of dimension $\text{Dim}(S)$, and sets of lower dimension will have Keisler measure 0.

Keisler measure behaves particularly well for NIP theories (theories without the independence property) so is orthogonal to the context of the present paper. For example, the authors show that if T is NIP, and μ is a Keisler measure on X , then there are boundedly many \sim_μ -classes of definable subsets of X , where $Y \sim_\mu Z$ whenever $\mu(Y \Delta Z) = 0$. They investigate existence and uniqueness of Keisler measures on groups.

Next, model-theoretic ideas of Euler characteristic were developed initially for o-minimal theories (see e.g. [17]) but more generally by Krajiček [35] and later Krajiček and Scanlon [36]. The latter define a *strong ordered Euler characteristic* on a structure M to be a function $\chi : \text{Def}(X) \rightarrow R$, where R is a partially ordered

ring, the image of χ takes values among the non-negative elements of R , and we have

- (a) $\chi(X) = \chi(Y)$ if X and Y are in definable bijection,
- (b) $\chi(X \times Y) = \chi(X) \cdot \chi(Y)$,
- (c) $\chi(X \cup Y) = \chi(X) + \chi(Y)$ if $X \cap Y = \emptyset$, and
- (d) $\chi(E) = c\chi(B)$ if $s : E \rightarrow B$ is a definable function and $c = \chi(f^{-1}(b))$ for each $b \in B$.

The Euler characteristic is *non-trivial* if $0 < 1$ in R and the image of χ is not just $\{0\}$.

Unlike for our notion of measure, there is no associated dimension, and indeed, the analogue of (c) for measure only holds if $\text{Dim}(X) = \text{Dim}(Y)$. However, under this definition very similar counting arguments are available. For example, Scanlon (see the Appendix of [40]) has shown that any field with strong ordered Euler characteristic is perfect and quasifinite, and these conclusions follow by the same argument for measurable fields.

Other questions about measure, and variations on measure are suggested by Hrushovski at the end of [29]. For example, there is a suggestion to consider finitely additive measures into $\mathbb{Q}[T]$, which could incorporate the current notion of dimension (exponent of leading term) and measure (coefficient of leading term).

References

- [1] M. Aschbacher, ‘On the maximal subgroups of the finite classical groups’, *Invent. Math.* 76 (1984), 469–514.
- [2] J. Ax, ‘The elementary theory of finite fields’, *Ann. Math.* 88 (1968), 239–271.
- [3] I. Ben-Yaacov, ‘On measurable first order theories and the continuous theory of measure algebras’, preprint.
- [4] I. Ben-Yaacov, I. Tomašić, F. O. Wagner, ‘Constructing an almost hyperdefinable group’, *J. Math. Logic*, 4 (2004), 181–212.
- [5] G. Bergman, H.W. Lenstra, ‘Subgroups close to normal subgroups’, *J. Algebra* 127 (1989), 80–97.
- [6] B. Bollobás, A. Thomason, ‘Graphs which contain all small graphs’, *Europ. J. Comb.* 2 (1981), 13–15.
- [7] B. Bollobás, *Random Graphs*, Academic Press, New York, 1985.
- [8] A. Borovik, G. Cherlin, ‘Permutation groups of finite Morley rank’, preprint.
- [9] R. Carter, *Simple groups of Lie type*, vol. XXVIII of *Pure and Applied Mathematics*, Wiley, 1972.
- [10] Z. Chatzidakis, L. van den Dries, A.J. Macintyre, ‘Definable sets over finite fields’, *J. Reine Angew. Math.* 427 (1992), 107–135.
- [11] Z. Chatzidakis, E. Hrushovski, ‘The model theory of difference fields’, *Trans. Amer. Math. Soc.* 351 (1999), 2997–3071.
- [12] Z. Chatzidakis, A. Pillay, ‘Generic structures and simple theories’, *Ann. Pure Appl. Logic* 95 (1998), 71–92.
- [13] Z. Chatzidakis, ‘Model theory of finite and pseudo-finite fields’, *Ann. Pure Appl. Logic* 88 (1997), 95–108.

- [14] G. Cherlin, E. Hrushovski, *Finite structures with few types*, Annals of Mathematics Studies No. 152, Princeton University Press, Princeton, 2003.
- [15] A. Chowdhury, B. Hart, Z. Sokolović, ‘Affine covers of Lie geometries and the amalgamation property’, Proc. London Math. Soc. (3) 85 (2002), 513–563.
- [16] L. van den Dries, ‘Algebraic theories with definable Skolem functions’, J. Symb. Logic 49 (1984), 625–629.
- [17] L. van den Dries, *Tame topology and o-minimal structures*, London Math. Soc Lecture Notes 248, Cambridge University Press, 1998.
- [18] R. Elwes, *Dimension and measure in finite first order structures*, PhD thesis, University of Leeds, 2005.
- [19] R. Elwes, M. Ryten, ‘Measurable groups of low dimension’, in preparation.
- [20] M. Fried, D. Haran, M. Jarden, ‘Effective counting of the points of definable sets over finite fields’, Isr. J. Math. 85 (1994), 103–133.
- [21] M. Fried, M. Jarden, *Field arithmetic*, 2nd Ed., Springer, Berlin, 2005.
- [22] R. L. Graham, J. H. Spencer, ‘A constructive solution to a tournament problem’, Canad. Math. Bull. 14 (1971), 45–48.
- [23] W. Hodges, *Model Theory*, Cambridge University Press, Cambridge, 1993.
- [24] E. Hrushovski, ‘Almost orthogonal regular types’, Ann. Pure Appl. Logic 45 (1989), 139–155.
- [25] E. Hrushovski, ‘Unimodular minimal structures’, J. London Math. Soc. (2) 46 (1992), 385–396.
- [26] E. Hrushovski, ‘A new strongly minimal set’, Ann. Pure Appl. Logic 62 (1993), 147–166.
- [27] E. Hrushovski, A. Pillay, ‘Definable subgroups of algebraic groups over finite fields’, J. Reine Angew. Math. 462 (1995), 69–91.
- [28] E. Hrushovski, ‘Simplicity and the Lascar group’, unpublished notes, 1997.
- [29] E. Hrushovski, ‘Pseudofinite fields and related structures’, in *Model theory and applications* (Eds. L. Bélair, Z. Chatzidakis, P. D’Aquino, D. Marker, M. Otero, F. Point, A. Wilkie), Quaderni di Matematica, vol. 11, Caserta, 2005, 151–212.
- [30] E. Hrushovski, ‘The first order theory of the Frobenius’, preprint.
- [31] E. Hrushovski, Y. Peterzil, A. Pillay, ‘Groups, measures, and the NIP’, preprint.
- [32] W.M. Kantor, M.W. Liebeck, H.D. Macpherson, ‘ \aleph_0 -categorical structures smoothly approximated by finite substructures’, Proc. London Math. Soc. (3) 59 (1989), 439–463.
- [33] H.J. Keisler, ‘Measures and forking’, Ann. Pure Appl. Logic 43 (1987), 119–169.
- [34] B. Kim, A. Pillay, ‘Simple theories’, Ann. Pure Appl. Logic 88 (1997), 149–164.

- [35] J. Krajiček, ‘Uniform families of polynomial equations over a finite field and structures admitting an Euler characteristic of definable sets’, *Proc. London Math. Soc.* (3) 81 (2000), 257–284.
- [36] J. Krajiček, T. Scanlon, ‘Combinatorics with definable sets: Grothendieck rings are Euler characteristics’, *Bull. Symb. Logic* 6 (2000), 311–330.
- [37] M.W. Liebeck, C.E. Praeger, J. Saxl, ‘On the O’Nan-Scott theorem for finite primitive permutation groups’, *J. Austral. Math. Soc.* 44 (1988), 389–396.
- [38] I.D. Macdonald, ‘Some explicit bounds in groups with finite derived subgroups’, *Proc. London Math. Soc.* (3) 11 (1961), 23–56.
- [39] H.D. Macpherson, A. Pillay, ‘Primitive permutation groups of finite Morley rank’, *Proc. London Math. Soc.* (3) 70 (1995), 481–504.
- [40] H.D. Macpherson, C. Steinhorn, ‘One-dimensional asymptotic classes of finite structures’, preprint.
- [41] H.D. Macpherson, K. Tent, ‘Stable pseudofinite groups’, *J. Alg.*, to appear.
- [42] A. Martin-Pizarro, A. Pillay, ‘Elliptic and hyperelliptic curves over supersimple fields’, *J. London Math. Soc.* (2) 69 (2004), 1–13.
- [43] A. Martin-Pizarro, F.O. Wagner, ‘Supersimplicity and quadratic extensions’, preprint.
- [44] B.H. Neumann, ‘Groups covered by permutable subsets’, *J. London Math. Soc.* 29 (1954), 236–248.
- [45] A. Pillay, *Geometric stability theory*, Oxford University Press, Oxford Logic Guides 32, 1996.
- [46] A. Pillay, T. Scanlon, F.O. Wagner, ‘Supersimple fields and division rings’, *Math. Research Letters* 5 (1998), 473–483.
- [47] F. Point, ‘Ultraproducts and Chevalley groups’, *Arch. Math. Logic* 38 (1999), 355–372.
- [48] M. Ryten, *Results around asymptotic and measurable groups*, PhD thesis, University of Leeds, in preparation.
- [49] M. Ryten, I. Tomašić, ‘ACFA and measurability’, *Selecta Mathematica*, to appear.
- [50] T. Scanlon, ‘Fields admitting nontrivial strong ordered Euler characteristics are quasifinite’, unpublished manuscript.
- [51] G. Schlichting, ‘Operationen mit periodischen Stabilisatoren’, *Arch. Math.* 34 (1980), 97–99.
- [52] W. Szmielew, ‘Elementary properties of abelian groups’, *Fund. Math.* 41 (1955), 203–271.
- [53] T. Szönyi, ‘Some applications of algebraic curves in finite geometry and combinatorics’, in *Surveys in Combinatorics* (Ed. R. Bailey), London Math. Society Lecture Notes 241, Cambridge University Press, Cambridge, 1997, pps. 197–236.
- [54] S. Thomas, *Classification theory of simple locally finite groups*, PhD thesis, University of London, 1983.

- [55] I. Tomašić, ‘Exponential sums in pseudofinite fields and applications’, *Illinois J. Math.* 48 (2004), 1235–1257.
- [56] I. Tomašić, ‘Independence, measure and pseudofinite fields’, *Selecta Math.*, to appear.
- [57] F.O. Wagner, *Simple theories*, Kluwer, Dordrecht, 2000.
- [58] J.S. Wilson, ‘On pseudofinite simple groups’, *J. London Math. Soc.* (2) 51 (1995), 471–490.